

**USDA-ARS/
U.S. Wheat and Barley Scab Initiative
FY07 Final Performance Report (approx. May 07 – April 08)
July 15, 2008**

Cover Page

PI:	Jose Costa
Institution:	University of Maryland
Address:	Dept. of Nat. Res. and Landscape Arch. 2102 Plant Sciences Building College Park, MD 20742-4452
E-mail:	costaj@umd.edu
Phone:	301-405-1317
Fax:	301-314-9308
Fiscal Year:	2007
USDA-ARS Agreement ID:	58-0790-5-078
USDA-ARS Agreement Title:	Screening and Developing Wheat Germplasm with Resistance to Scab.
FY07 ARS Award Amount:	\$ 27,661

USWBSI Individual Project(s)

USWBSI Research Area *	Project Title	ARS Adjusted Award Amount
HGG	Sources of Variation for DON Concentration in Moderately Resistant SRWW.	\$4,917
VDUN	Developing and Screening Wheat Germplasm with Resistance to Scab.	\$ 22,744
	Total Award Amount	\$ 27,661

Principal Investigator

Date

* CBCC – Chemical, Biological & Cultural Control
EEDF – Etiology, Epidemiology & Disease Forecasting
FSTU – Food Safety, Toxicology, & Utilization of Mycotoxin-contaminated Grain
GET – Genetic Engineering & Transformation
HGR – Host Genetics Resources
HGG – Host Genetics & Genomics
IIR – Integrated/Interdisciplinary Research
PGG – Pathogen Genetics & Genomics
VDUN – Variety Development & Uniform Nurseries

Project 1: *Sources of Variation for DON Concentration in Moderately Resistant SRWW.*

1. What major problem or issue is being resolved and how are you resolving it?

In collaboration with Dr. Caly Sneller (Ohio State University) we are attempting to understand variation for DON accumulation among moderately resistant lines to efficiently obtain cultivars with low scab index and low DON required by the wheat industry.

We are testing a large set of soft red winter wheat breeding lines with moderate resistance to FHB. During the 2007-2008 we grew these lines, their parents, and checks, and rated them for FHB in June 2007 at Feekes' GS 11.2. We collected grain samples that will be tested for DON in the fall of 2008.

**2. List the most important accomplishment and its impact (how is it being used?).
Complete all three sections (repeat sections for each major accomplishment):**

At this time we are processing grain samples harvested in June 2008 for DON testing. No major accomplishment has been achieved at this time. It is the first season of testing in MD for this project.

Project 2: *Developing and Screening Wheat Germplasm with Resistance to Scab.***1. What major problem or issue is being resolved and how are you resolving it?**

The major problem being addressed is the need to develop rapidly and effectively host resistance to scab (*Fusarium Head Blight*) from exotic sources into adapted soft red winter wheat (SRWW) germplasm. The approach to address this problem is to backcross, and to three-way cross the Sumai 3 allele and other exotic resistance alleles into adapted SRWW lines and varieties. Marker-assisted selection (MAS) is being used to rapidly incorporate Sumai 3 (from Ning7840) resistance into SRWW lines such as McCormick, that have wide adaptation in the Southern and Eastern US wheat growing regions and moderate resistance to scab. Over 400 Backcross 2 F1 seedlings were screened for SSR markers at the USDA in Raleigh (NC) in collaboration with Dr. Gina Brown-Guedira at that National Genotyping Center. Of these, a single BC2F1 plant was selected that had Ning7840 alleles at the 3BS, 5A and 2DL genomic regions. 340 F2 progenies were further screened in 2007, selected homozygous BC2F3s were planted in the field in the fall of 2007 in our inoculated nursery. Incidence and severity of scab were scored in the spring of 2008 and these will be tested for DON in the fall of 2008.

Additionally, selected BC-1F1s were crossed with the wheat cultivar SS 8641 that has wide adaptation as well as leaf and stripe rust resistance. The BC1 F1 seeds were screened with markers in 2007-2008 and the BC1-F2s selected progenies were advanced in the growth chamber. Selected homozygous BC1F3s for the 3BS, 5A and 2DL genomic regions, will be planted in the field in the fall of 2008.

Furthermore, screening of MD (University of Maryland) wheat advanced lines and check varieties was conducted under field conditions in an inoculated nursery at Salisbury (MD). Conditions favorable for disease development were aided with daily misting before and during wheat flowering. The scab inoculum was scabby corn grain spread in the field a month before flowering. The Southern wheat scab and Northern Uniform Scab Screening nurseries that include new experimental lines were also screened for resistance at Salisbury (MD) with artificial inoculation and misting. Data for all nurseries was obtained for scab incidence, scab severity, *Fusarium* damaged kernels, seed weight, plant height, heading date, and DON levels.

2. List the most important accomplishment and its impact (how is it being used?).

Complete all three sections (repeat sections for each major accomplishment):

Accomplishment: Incorporation of the 3BS, 5A and 2DL quantitative trait loci (QTL) of resistance to scab from Sumai3 into adapted soft red winter wheat germplasm.

Impact: the availability of these germplasm with resistance will reduce scab negative effects in years favorable to scab development.

As a result of that accomplishment, what does your particular clientele, the scientific community, and agriculture as a whole have now that they didn't have before?

Plant breeders will have homozygous F4 seed for the 3 QTL of adapted soft red winter wheat germplasm available in the McCormick and SS8641 background in the fall of 2008 for crossing.

Include below a list of the publications, presentations, peer-reviewed articles, and non-peer reviewed articles written about your work that resulted from all of the projects included in the grant. Please reference each item using an accepted journal format. If you need more space, continue the list on the next page.

Costa, J. M., Al-Tukhaim, L., Gal-Edd, N., Ku, A., Wenger, E., Brown-Guedira, G., and Van Sanford, D. 2007. Development of Scab Resistant Soft Red Winter Wheat Germplasm (SRWW) Using Marker-Assisted Selection (MAS). Proceedings of the 2007 National Fusarium Head Blight Forum p. 175. Kansas City, MI.

Presentations:

Costa, J.M. 2007. Use of DNA markers for developing head blight resistant wheat genotypes. 2007 Argentina Symposium on wheat diseases. University of La Plata, Argentina. 11/23/2007.

Costa, J.M. 2008. Marker-assisted selection for head blight resistance in winter wheat. University of Puerto Rico. Mayaguez, PR. 1/17/2008.