

**USDA-ARS/
U.S. Wheat and Barley Scab Initiative
FY10 Final Performance Report
July 15, 2011**

Cover Page

PI:	Elias Elias
Institution:	North Dakota State University
Address:	Department of Plant Sciences NDSU Dept. # 7670 PO Box 6050 Fargo, ND 58108-6050
E-mail:	elias.elias@ndsu.edu
Phone:	701-231-8159
Fax:	701-231-8474
Fiscal Year:	FY10
USDA-ARS Agreement ID:	59-0206-9-061
USDA-ARS Agreement Title:	Identify and Develop Durum Wheat Resistant to Fusarium Head Blight.
FY10 USDA-ARS Award Amount:	\$ 128,780

USWBSI Individual Project(s)

USWBSI Research Category*	Project Title	ARS Award Amount
DUR-CP	Develop Durum Wheat Resistant to Fusarium Head Blight.	\$ 92,683
DUR-CP	Identify and Develop Durum Wheat Resistant to Fusarium Head Blight.	\$ 36,097
	Total ARS Award Amount	\$ 128,780

Principal Investigator

Date

* MGMT – FHB Management
 FSTU – Food Safety, Toxicology, & Utilization of Mycotoxin-contaminated Grain
 GDER – Gene Discovery & Engineering Resistance
 PBG – Pathogen Biology & Genetics
 BAR-CP – Barley Coordinated Project
 DUR-CP – Durum Coordinated Project
 HWW-CP – Hard Winter Wheat Coordinated Project
 VDHR – Variety Development & Uniform Nurseries – Sub categories are below:
 SPR – Spring Wheat Region
 NWW – Northern Soft Winter Wheat Region
 SWW – Southern Soft Red Winter Wheat Region

Project 1: *Develop Durum Wheat Resistant to Fusarium Head Blight.*

1. What major problem or issue is being resolved relevant to Fusarium head blight (scab) and how are you resolving it?

Fusarium head blight (FHB) is caused by the fungus *Fusarium graminearum* Schwabe (telomorph *Gibberella zeae* (Schwein.) Petch., which has been seriously attacking durum wheat. Since 1993, it is estimated that FHB has cost over \$3 billion in direct and indirect losses in North Dakota. Although fungicides may reduce FHB, using genetic resistance is the most environmentally safe and economical way to control the disease. The objective of this project is to incorporate identified sources of resistance into the currently susceptible durum wheat germplasm in order to develop resistant cultivars.

2. List the most important accomplishment and its impact (i.e. how is it being used) to minimize the threat of Fusarium head blight or to reduce mycotoxins. Complete both sections (repeat sections for each major accomplishment):

Accomplishment:

The cultivar Divide that has some level of resistance to FHB is gaining acreages in North Dakota. The acreage of Divide increased from 20.6 % in 2009 to 26.6% in 2010 of the total durum acreage in North Dakota.

- Sumai , Wangshuibai, and Tunisian sources of resistance:
 - 2 lines were evaluated in the Uniform Regional Nursery
 - 14 lines were evaluated in the Elite Advanced Yield Trial
 - 188 lines were evaluated in the Advanced Yield Trials
 - 640 lines were evaluated in the Preliminary Yield Trials
 - 50 populations were screened in the field and greenhouses
 - 71 new populations were developed

Impact:

The above developed material is the only known improved durum germplasm with Fusarium head blight resistance. This germplasm is vital for the survival of the Midwest durum producers. Since the Midwest produces over 65% of the US durum, this germplasm has a major impact on the pasta industry and the US economy. Divide, based on its FHB resistance and yield advantage and current grown acreage will generate additional millions of dollars into the economy.

Project 2: Identify and Develop Durum Wheat Resistant to Fusarium Head Blight.

1. What major problem or issue is being resolved relevant to Fusarium head blight (scab) and how are you resolving it?

Durum Wheat is very susceptible to Fusarium head blight (FHB) caused by the fungus *Fusarium graminearum* Schwabe (teleomorph *Gibberella Zeae* (Schw.) Petch. Sources of resistance to FHB in durum wheat that are equivalent to the Chinese spring wheat Sumai 3 are not available yet. Our objective is to identify sources of resistance that can be utilized by durum plant breeders to develop FHB resistant cultivars. To date we have screened all the durum wheat accessions in the National small grain Collection, Aberdeen, ID. There are 15,000 durum wheat accessions at the International Center of Agricultural Research in the Dry Areas (ICARDA) and International Maize and Wheat Improvement Center (CIMMYT). We are in the process of evaluating these accessions in field nurseries in China and greenhouses in North Dakota.

2. List the most important accomplishment and its impact (i.e. how is it being used) to minimize the threat of Fusarium head blight or to reduce mycotoxins. Complete both sections (repeat sections for each major accomplishment):

Accomplishment:

- To date we have evaluated 3,526 accessions from ICARDA. After several evaluations in the field and greenhouses four accessions maintained disease severity less than 30%. Twenty seven populations were developed from crossing the four accessions with adapted germplasm.
- Nine hundred- eighty-five new accessions were sent to China for evaluation. These accessions were held in quarantines in China and therefore will be evaluated in 2011.
- Fifteen selected ICARDA accessions were reevaluated in the spring 2010 greenhouse.
- Thirty-seven lines from crosses with Tunisian lines were evaluated in Advanced Yield Trial.
- Nine lines from crosses with Tunisian lines were evaluated in the Elite Advanced Yield Trial.
- Two hundred-twenty lines from crosses with CIMMYT lines were evaluated in preliminary yield trials.
- Ten populations were developed from crossing adapted germplasm with Tunisian 7.
- Twenty populations were developed from crossing adapted germplasm with *Triticum dicoccum*.

Impact:

Any resistant germplasm that is identified above could potentially lead to the development of FHB resistant durum cultivars. Resistant durum cultivars will generate millions of dollars into the farm economy in the Midwest and will insure the stability of the durum industry in the United States.

FY10 (approx. May 10 – May 11)
PI: Elias, Elias
USDA-ARS Agreement #: 59-0206-9-061

FY10 Final Performance Report

Include below a list of all germplasm or cultivars released with full or partial support of the USWBSI. List the release notice or publication. Briefly describe the level of FHB resistance.

None

Include below a list of the publications, presentations, peer-reviewed articles, and non-peer reviewed articles written about your work that resulted from all of the projects included in the grant. Please reference each item using an accepted journal format. If you need more space, continue the list on the next page.

None